

Write the Date received in the box next to the vaccine once your child has recieved it.

НерВ	Hepatitis B (this dose may be administered anywhere from 6-18 months. Talk to your provider for more information.)	(Dose 3 of 3)	Date received
RV*	Rotavirus	(Dose 3 of 3)	Date received
DTaP	Diphtheria, Pertussis, & Tetanus	(Dose 3 of 5)	Date received
Hib*	Haemophilus influenzae type b	(Dose 3 of 4)	Date received
PCV15, PCV20	Pneumococcal disease	(Dose 3 of 4)	Date received
IPV	Polio (This dose may be administered anywhere from 6 -18 months. Talk to your provider for more information.)	(Dose 3 of 4)	Date received
COVID-19**	Coronavirus disease 2019 (1 or more doses of the updated (2023–2024 Formula) vaccine is recommended. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
Flu†	Influenza (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
RSV*	Respiratory Syncytial Virus (This dose can be administered anywhere from birth-19 months. Talk to your doctor or health care provider for more information.)	(Dose 1 of 1)	Date received

FOOTNOTES

RV* Hib*

Administering a third dose at age 6 months depends on the brand of Hib or rotavirus vaccine used for previous dose.

COVID-19**

Number of doses recommended depends on your child's age and type of COVID-19 vaccine used.

Flu[†]

Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group. The flu shot should be received yearly.

RSV*

Your child might need an RSV vaccine. Talk to your doctor or health care provider for more information.



DISCLAIMER: Dosages and Vaccinations required are as presented on the CDC's recommended infant vaccination schedule. Contact your health care provider or visit https://www.cdc.gov/vaccines for more information.

Last Updated: April 2024