

2024 Recommended Immunizations for Children Aged

12 Months

Write the Date received in the box next to the vaccine once your child has received it.

HepB	Hepatitis B (This dose may be administered anywhere from 6–18 months. Talk to your provider for more information.)	(Dose 3 of 3)	Date received
Hib	Haemophilus influenzae type b (This dose may be administered anywhere from 12–15 months. Talk to your provider for more information.)	(Dose 4 of 4)	Date received
PCV15, PCV20	Pneumococcal disease (This dose may be administered anywhere from 12–15 months. Talk to your provider for more information.)	(Dose 4 of 4)	Date received
IPV	Polio (This dose may be administered anywhere from 6–18 months. Talk to your provider for more information.)	(Dose 3 of 4)	Date received
COVID-19**	Coronavirus disease 2019 (1 or more doses of the updated (2023–2024 Formula) vaccine is recommended. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
Flu†	Influenza (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
MMR	Measles, Mumps, & Rubella (This dose may be administered anywhere from 6–18 months. Talk to your provider for more information.)	(Dose 1 of 2)	Date received
Varicella	Chickenpox (This dose may be administered anywhere from 12–15 months. Talk to your provider for more information.)	(Dose 1 of 2)	Date received
HepA‡	Hepatitis A	(Dose 1 of 2)	Date received
RSV*	Respiratory Syncytial Virus (This dose can be administered anywhere from birth–19 months. Talk to your doctor or health care provider for more information.)	(Dose 1 of 1)	Date received

FOOTNOTES

COVID-19**

Number of doses recommended depends on your child's age and type of COVID-19 vaccine used.

Flu†

Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group. The flu shot should be received yearly.

HepA‡

Two doses of Hep A vaccine are needed for lasting protection. The 2 doses should be given between age 12 and 23 months. Both doses should be separated by at least 6 months. Children 2 years and older who have not received 2 doses of Hep A should complete the series.

RSV*

Your child might need an RSV vaccine. Talk to your doctor or health care provider for more information.



DISCLAIMER: Dosages and Vaccinations required are as presented on the CDC's recommended infant vaccination schedule. Contact your health care provider or visit <https://www.cdc.gov/vaccines> for more information.

Last Updated: April 2024