

## Recommended Immunizations for Children Aged

# 12 Months

Write the Date received in the box next to the vaccine once your child has received it.

<b>HepB</b>	<b>Hepatitis B</b> (this dose may be administered anywhere from 6-18 months. Talk to your provider for more information.)	(Dose 3 of 3)	Date received
<b>Hib</b>	<b>Haemophilus influenzae type b</b> (This dose may be administered anywhere from 12-15 months. Talk to your provider for more information.)	(Dose 4 of 4)	Date received
<b>PCV15, PCV20</b>	<b>Pneumococcal disease</b> (This dose may be administered anywhere from 12-15 months. Talk to your provider for more information.)	(Dose 4 of 4)	Date received
<b>IPV</b>	<b>Polio</b> (This dose may be administered anywhere from 6-18 months. Talk to your provider for more information.)	(Dose 3 of 4)	Date received
<b>COVID-19**</b>	<b>Coronavirus disease 2019</b> (1 or more doses of the updated (2024–2025 Formula) vaccine is recommended. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
<b>Flu†</b>	<b>Influenza</b> (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
<b>MMR</b>	<b>Measles, Mumps, &amp; Rubella</b> (This dose may be administered anywhere from 6-18 months. Talk to your provider for more information.)	(Dose 1 of 2)	Date received
<b>Varicella</b>	<b>Chickenpox</b> (This dose may be administered anywhere from 12-15 months. Talk to your provider for more information.)	(Dose 1 of 2)	Date received
<b>HepA‡</b>	<b>Hepatitis A</b>	(Dose 1 of 2)	Date received
<b>RSV*</b>	<b>Respiratory Syncytial Virus</b> (This dose can be administered anywhere from birth-19 months. Talk to your doctor or health care provider for more information.)	(Dose 1 of 1)	Date received

### FOOTNOTES

#### COVID-19\*\*

Number of doses recommended depends on your child's age and type of COVID-19 vaccine used.

#### Flu†

Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group. The flu shot should be received yearly.

#### HepA‡

Two doses of Hep A vaccine are needed for lasting protection. The 2 doses should be given between age 12 and 23 months. Both doses should be separated by at least 6 months. Children 2 years and older who have not received 2 doses of Hep A should complete the series.

#### RSV\*

Your child might need an RSV vaccine. Talk to your doctor or health care provider for more information.



**DISCLAIMER:** Dosages and Vaccinations required are as presented on the CDC's recommended infant vaccination schedule. Contact your health care provider or visit <https://www.cdc.gov/vaccines> for more information.

Last Updated: February 2025