

Recommended Immunizations for Children Aged

4-6 Years

Write the Date received in the box next to the vaccine once your child has received it.

DTaP	Diphtheria, Pertussis, & Tetanus	(Dose 5 of 5)	Date received
IPV	Polio	(Dose 4 of 4)	Date received
COVID-19**	Coronavirus disease 2019 (1 or more doses of the updated (2024–2025 Formula) vaccine is recommended. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
Flu†	Influenza (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
MMR	Measles, Mumps, & Rubella	(Dose 2 of 2)	Date received
Varicella	Chickenpox	(Dose 2 of 2)	Date received

FOOTNOTES

COVID-19**

Number of doses recommended depends on your child's age and type of COVID-19 vaccine used.

Flu†

Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group. The flu shot should be received yearly.



DISCLAIMER: Dosages and Vaccinations required are as presented on the CDC's recommended infant vaccination schedule. Contact your health care provider or visit <https://www.cdc.gov/vaccines> for more information.

Last Updated: February 2025