

Recommended Immunizations for Children Aged

12 Months

Write the Date received in the box next to the vaccine once your child has received it.

| | | | |
|---------------------|--|---------------|---------------|
| HepB | Hepatitis B (this dose may be administered anywhere from 6-18 months. Talk to your provider for more information.) | (Dose 3 of 3) | Date received |
| Hib | Haemophilus influenzae type b (This dose may be administered anywhere from 12-15 months. Talk to your provider for more information.) | (Dose 4 of 4) | Date received |
| PCV15, PCV20 | Pneumococcal disease (This dose may be administered anywhere from 12-15 months. Talk to your provider for more information.) | (Dose 4 of 4) | Date received |
| IPV | Polio (This dose may be administered anywhere from 6-18 months. Talk to your provider for more information.) | (Dose 3 of 4) | Date received |
| COVID-19** | Coronavirus disease 2019 (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.) | (Dose 1 of 1) | Date received |
| Flu† | Influenza (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.) | (Dose 1 of 1) | Date received |
| MMR | Measles, Mumps, & Rubella (This dose may be administered anywhere from 12 to 15 months. Talk to your provider for more information.) | (Dose 1 of 2) | Date received |
| Varicella | Chickenpox (This dose may be administered anywhere from 12-15 months. Talk to your provider for more information.) | (Dose 1 of 2) | Date received |
| HepA‡ | Hepatitis A (This dose may be administered anywhere from 12-17 months. Talk to your provider for more information.) | (Dose 1 of 2) | Date received |
| RSV* | Respiratory Syncytial Virus (This dose can be administered anywhere from birth-19 months. Talk to your doctor or health care provider for more information.) | (Dose 1 of 1) | Date received |

FOOTNOTES

COVID-19**

Annual COVID-19 vaccination is recommended for children 6-23 months and yearly for children 2 years and older in specific groups or if parents choose protection. The number of doses depends on your child's age and vaccine type.

Flu†

Two doses given at least 4 weeks apart are recommended for children aged 6 months through 8 years who are getting an influenza (flu) vaccine for the first time, and for some other children in this age group. The flu shot should be received yearly.

HepA‡

Two doses of Hep A vaccine are needed for lasting protection. The two doses should be given between the ages of 12 and 23 months. Both doses should be separated by at least 6 months. Children 2 years and older who have not received 2 doses of Hep A should complete the series.

RSV*

Your child might need an RSV vaccine. Talk to your doctor or health care provider for more information.



DISCLAIMER: Dosages and Vaccinations required are as presented on the Illinois Department of Public Health's recommended infant vaccination schedule. Contact your health care provider or visit <https://dph.illinois.gov/topics-services/prevention-wellness/immunization.html> for more information.

Last Updated: February 2026