

Recommended Immunizations for Children Aged

15 Months

Write the Date received in the box next to the vaccine once your child has received it.

HepB	Hepatitis B (This dose may be administered anywhere from 6-18 months. Talk to your provider for more information.)	(Dose 3 of 3)	Date received
DTaP	Diphtheria, Pertussis, & Tetanus (This dose may be administered anywhere from 15-18 months. Talk to your provider for more information.)	(Dose 4 of 5)	Date received
Hib	Haemophilus influenzae type b (This dose may be administered anywhere from 12-15 months. Talk to your provider for more information.)	(Dose 4 of 4)	Date received
PCV15, PCV20	Pneumococcal disease (This dose may be administered anywhere from 12-15 months. Talk to your provider for more information.)	(Dose 4 of 4)	Date received
IPV	Polio (This dose may be administered anywhere from 6-18 months. Talk to your provider for more information.)	(Dose 3 of 4)	Date received
COVID-19**	Coronavirus disease 2019 (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
Flu†	Influenza (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
MMR	Measles, Mumps, & Rubella (This dose may be administered anywhere from 12-18 months. Talk to your provider for more information.)	(Dose 1 of 2)	Date received
Varicella	Chickenpox (This dose may be administered anywhere from 12-15 months. Talk to your provider for more information.)	(Dose 1 of 2)	Date received
RSV*	Respiratory Syncytial Virus (This dose can be administered anywhere from birth-19 months. Talk to your doctor or health care provider for more information.)	(Dose 1 of 1)	Date received
HepA‡	Hepatitis A (This dose may be administered anywhere from 12-17 months. Talk to your provider for more information.)	(Dose 1 of 2)	Date received

FOOTNOTES

COVID-19**

Annual COVID-19 vaccination is recommended for children ages 6-23 months. For children ages 2 years and older, the vaccine is also recommended yearly if they fall under specific risk groups or if the parents choose to provide protection against COVID-19.

Number of doses depends on your child's age and vaccine type.

Flu†

Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group. The flu shot should be received yearly.

RSV*

Your child might need an RSV vaccine. Talk to your doctor or health care provider for more information.

HepA‡

Two doses of Hep A vaccine are needed for lasting protection. The 2 doses should be given between age 12 and 23 months. Both doses should be separated by at least 6 months. Children 2 years and older who have not received 2 doses of Hep A should complete the series.



DISCLAIMER: Dosages and Vaccinations required are as presented on the Illinois Department of Public Health's recommended infant vaccination schedule. Contact your health care provider or visit <https://dph.illinois.gov/topics-services/prevention-wellness/immunization.html> for more information.

Last Updated: February 2026