

## Recommended Immunizations for Children Aged

# 18 Months

Write the Date received in the box next to the vaccine once your child has received it.

<b>HepB</b>	<b>Hepatitis B</b> (This dose may be administered anywhere from 6-18 months. Talk to your provider for more information.)	(Dose 3 of 3)	Date received
<b>DTaP</b>	<b>Diphtheria, Pertussis, &amp; Tetanus</b> (This dose may be administered anywhere from 15-18 months. Talk to your provider for more information.)	(Dose 4 of 5)	Date received
<b>IPV</b>	<b>Polio</b> (This dose may be administered anywhere from 6-18 months. Talk to your provider for more information.)	(Dose 3 of 4)	Date received
<b>COVID-19**</b>	<b>Coronavirus disease 2019</b> (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
<b>Flu†</b>	<b>Influenza</b> (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
<b>HepA‡</b>	<b>Hepatitis A</b> (This dose may be administered anywhere from 18-23 months. Talk to your provider for more information.)	(Dose 2 of 2)	Date received
<b>RSV*</b>	<b>Respiratory Syncytial Virus</b> (This dose can be administered anywhere from birth-19 months. Talk to your doctor or health care provider for more information.)	(Dose 1 of 1)	Date received

## FOOTNOTES

### COVID-19\*\*

Annual COVID-19 vaccination is recommended for children 6-23 months, and yearly for children 2 years and older in specific groups or if parents choose protection. Number of doses depends on your child's age and vaccine type.

### Flu†

Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group. The flu shot should be received yearly.

### HepA‡

Two doses of Hep A vaccine are needed for lasting protection. The 2 doses should be given between age 12 and 23 months. Both doses should be separated by at least 6 months. Children 2 years and older who have not received 2 doses of Hep A should complete the series.

### RSV\*

Your child might need an RSV vaccine. Talk to your doctor or health care provider for more information.



**DISCLAIMER:** Dosages and Vaccinations required are as presented on the Illinois Department of Public Health's recommended infant vaccination schedule. Contact your health care provider or visit <https://dph.illinois.gov/topics-services/prevention-wellness/immunization.html> for more information.

Last Updated: February 2026