

Recommended Immunizations for Children Aged

4-6 Years

Write the Date received in the box next to the vaccine once your child has received it.

DTaP	Diphtheria, Pertussis, & Tetanus	(Dose 5 of 5)	Date received
IPV	Polio	(Dose 4 of 4)	Date received
COVID-19**	Coronavirus disease 2019 (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
Flu†	Influenza (These vaccinations are given once a year, starting at 6 months. Talk to your provider for more information.)	(Dose 1 of 1)	Date received
MMR	Measles, Mumps, & Rubella	(Dose 2 of 2)	Date received
Varicella	Chickenpox	(Dose 2 of 2)	Date received

FOOTNOTES

COVID-19**

Annual COVID-19 vaccination is recommended for children 6-23 months, and yearly for children 2 years and older in specific groups or if parents choose protection. Number of doses depends on your child's age and vaccine type.

Flu†

Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group. The flu shot should be received yearly.



DISCLAIMER: Dosages and Vaccinations required are as presented on the Illinois Department of Public Health's recommended infant vaccination schedule. Contact your health care provider or visit <https://dph.illinois.gov/topics-services/prevention-wellness/immunization.html> for more information.

Last Updated: February 2026